wherein M is Fe[II], Fe[III], Co[I], Co[II], Mn[I], Mn[II], Mn[II], Mn[IV], Ru[II], Ru[III] or Ru[IV]; X represents an atom or group covalently or ionically bonded to the transition metal M;

R is independently selected from hydrogen, halogen, hydrocarbyl, substituted hydrocarbyl, heterohydrocarbyl or substituted heterohydrocarbyl;

Z is a bridging group comprising a donor atom of N, P or S or alternatively is a neutral group comprising a C_{1-4} alkylene group, a silyl or germyl group; and n = an integer to satisfy the valency of M, and

(2) an activating quantity of an activator compound.

3

FINNEGAN HENDERSON FARABOW GARRETT & DUNNERLLP

- 40. (New) A polymerization catalyst comprising
- (1) a transition metal complex having the formula:

wherein M is Fe[II], Fe[III], Co[I], Co[II], Mn[II], Mn[II], Mn[III], Mn[IV], Ru[III] or Ru[IV]; X represents an atom or group covalently or ionically bonded to the transition metal M; Z is a bridging group comprising a donor atom of N, P or S or alternatively is a neutral group comprising a C_{1-4} alkylene group, a silyl or germyl group, R^1-R^{14} are independently selected from hydrogen, halogen, hydrocarbyl, substituted hydrocarbyl, heterohydrocarbyl or substituted heterohydrocarbyl; and n = an integer to satisfy the valency of M, and

(2) an activating quantity of an activator compound.

3

FINNEGAN
HENDERSON
FARABOW
GARRETT &
DUNNER

- 41. (New) A polymerization catalyst comprising
- (1) a complex having the formula

$$R^{12}$$

$$R^{11}$$

$$R^{12}$$

$$R^{13}$$

$$R^{14}$$

$$R^{10}$$

$$R^{10}$$

$$R^{10}$$

$$R^{10}$$

$$R^{10}$$

wherein M is Fe[II], Fe[III], Ni[II], Co[I], Co[II], Co[III], Mn[I], Mn[II], Mn[III], Mn[IV], Ru[II], Ru[IV], Pd[II], V[III], V[IV] or V[V];

X represents an atom or group covalently or ionically bonded to the transition metal M; Z is a bridging group comprising a donor atom of N, P or S or alternatively is a neutral group comprising a C_{1-4} alkylene group, a silyl or germyl group,

 R^{1} - R^{14} are independently selected from hydrogen, halogen, hydrocarbyl, substituted hydrocarbyl, heterohydrocarbyl, or substituted heterohydrocarbyl, and at least one of R^{1} - R^{10} contains two or more carbon atoms; and n = 1 an integer to satisfy the valency of M, and

- (2) an activating quantity of an activator compound.
- 42. (New) The polymerization catalyst of claim 40 or 41 wherein R¹¹-R¹⁴ are phenyl, alkyl or cycloalkyl.

3

FINNEGAN HENDERSON FARABOW GARRETT & DUNNER LLP

- 43. (New) The polymerization catalyst of claim 39, 40, or 41 wherein the bridging group Z is $-CH_2$ or a donor atom N.
 - 44. (New) The polymerization catalyst of claim 40 or 41 having the formula:

wherein R is hydrogen or hydrocarbyl.

- 45. (New) The polymerization catalyst of claim 39, 40, or 41 wherein the metal M is Fe or Co.
- 46. (New) The polymerization catalyst of claim 39, 40, or 41 wherein the Group X is chloride.
- 47. (New) The polymerization catalyst of claim 39, 40, or 41 wherein the activator compound is an organoaluminum compound or a hydrocarbylboron compound.
- 48. (New) The polymerization catalyst of claim 39, 40, or 41 further comprising a neutral Lewis base.
- 49. (New) The polymerization catalyst of claim 39, 40, or 41 further comprising a support.



FINNEGAN HENDERSON FARABOW GARRETT & DUNNERLLP